

Luis de Sarro 501 (B1838DQK) Luis Guillón, Buenos Aires, Argentina. (011) 4367-6100 l syntex@syntexar.com www **syntexar** com tecnologías para la **vida**

<u>GONASYN</u>

Introduction:

GONASYN is an injectable solution containing Gonadorelin Acetate, a synthetic analogue of hypothalamic releasing factor (GnRH).

Action:

Gonadorelin acetate stimulates the synthesis and release of follicle stimulating hormone (FSH) and luteinizing hormone (LH) from pituitary gland.

Formula:

Gonadorelin acetate50 µg Excipients e.q.1 mL

Species which the veterinary medicinal product is intended for: Bovine, Ovine, Caprine, Equine, Rabbits and Canine.

Therapeutic indications:

Bovine:

- Treatment of ovarian follicular cysts.
- Oestrus induction in anoestrus cows.
- Ovulation synchronization.
- Fertility improvement.

Ovine & Caprine:

• Ovulation induction.

<u>Equine:</u>

• Ovulation induction.

Rabbits:

• Ovulation induction.

Canine:

- Criptorquidism.
- Treatment of poly-cysts ovaries.



Luis de Sarro 501 (B1838DQK) Luis Guillón, Buenos Aires, Argentina. (011) 4367-6100 l syntex@syntexar.com www **syntexar** com tecnologías para la **vida**

Dosage:

<u>Bovine</u>: 100 μg/animal (2 ml) Ovine & Caprine: 100 μg/animal (2 ml)

Equine: 100 µg/animal (2 ml)

The administration has to be carried out only when gynaecological testing diagnoses an ovary follicle >35 mm, in case of ovulation induction failure the process can be repeated after a period longer than 24-36 hs.

<u>Rabbits</u>: 10-20 µg/animal (0.2-0.4 ml)

Canine: Criptorquidism: 50-100 µg/animal (1-2 ml)

In case of not observing the desired effect the dose can be repeated after 4-6 days.

Treatment of poly-cysts ovaries: 3.3 µg/kg (6.6 ml). Administer daily during 3 days.

Route of administration:

Intramuscular or intravenous injection

Toxicity data:

The product is not toxic at recommended doses.

There have been no cases of acute toxicity at recommended dosage.

The internationally established criteria consider these products as safe.

DL₅₀ (rats) SC = >2 g/kg DL₅₀ (rats) IV = >203 mg/kg DL₅₀ (rats) Oral = >3 g/kg DL₅₀ (mouse) Oral = >4 g/kg DL₅₀ (mouse) SC = >3 g/kg DL₅₀ (mouse) IV = >416 mg/kg

<u>Pharmaceutical form:</u> Injectable aqueous solution

Commercial presentations:

-Vial containing 20 ml of Gonadorelin Acetate 50 $\mu\text{g/ml}$

-Vial containing 100 ml of Gonadorelin Acetate 50 μ g/ml



<u>Withdrawal period</u>: MEAT: 0 days. MILK: 0 days Luis de Sarro 501 (B1838DQK) Luis Guillón, Buenos Aires, Argentina. (011) 4367-6100 I syntex@syntexar.com www **syntexar** com tecnologías para la **vida**

Precautions:

- Store below 25°C (air conditioning), protected from light. Do not freeze.
- In transport, should be kept in a cool, dry place and protected from light.
- No special care is required in the destruction of the container.

PRODUCT USE FOR REPRODUCTION MANAGEMENT

VETERINARY USE ONLY

KEEP OUT OF REACH OF CHILDREN

Manufactured by:

SYNTEX S.A.

Luis de Sarro 501 – (B1838DQK) Luis Guillón - Bs. As. – Argentina Telephone/fax: +54 (011) 4367-6100 E-mail: <u>syntex@syntexar.com</u>